

IN SEARCH OF AN EVIDENCE-BASED PRACTICE TO REDUCE DURATION OF UNTREATED PSYCHOSIS (DUP)

Wasef Atiya¹, Serena Chaudhry², Ashley Weiss², Robert Roy², Sydney Long³

¹ BA, MD/MBA Candidate at Tulane School of Medicine and Tulane Freeman School of Business, ² Early Psychosis Intervention Clinic New Orleans (EPIC-NOLA), Tulane University School of Medicine Department of Psychiatry and Behavioral Sciences, ³ BS, Candidate in Neuroscience in the School of Science and Engineering at Tulane University

INTRODUCTION

- Duration of untreated psychosis (DUP) is important in predicting outcomes for individuals with first-episode psychosis.
- DUP has two components: "supply" (duration from symptom onset to Coordinated Specialty Care clinic entry) and "demand" (duration from symptom onset to first antipsychotic medication).
- The Treatment and Intervention in Psychosis (TIPS) clinic and Mind Map program both demonstrated reduction in community DUP through psychosis early detection campaigns
- The Early Psychosis Intervention Clinic-New Orleans (EPIC-NOLA) aims to reduce DUP for clinic referrals by implementing a psychosis early detection campaign.

METHODS

- CALM was re-launched with an approach similar to Mind Map, with a 3-pronged approach to brand building:
 - Mass Transit Marketing
 - Digital Marketing
 - Local community engagement activities
- Descriptive data was collected for each method of engagement
- Website traffic, social media analytics, and reports from partner agencies were used to assess visibility of CALM campaign.
- Comparative analysis was used to examine DUP supply and demand before and after the re-launch of the CALM campaign.

RESULTS

DUP Component	Before CALM Launch	After CALM Launch
Total	9.85 months	5.95 months
Demand	6.29 months	2.25 months
Supply	4.21 months	3.06 months

Website Performance	Result
Website Traffic Increase	24%
Quizzes Completed	777
Listserv Subscribers	35

Social Media Analytics	June 15, 2022	June 15, 2023	Percent Change
Social Impressions	2,859	205,842	+7,079%
Social Engagement	278	11,616	+4,063.4%
Social Media Followers	1290	1517	+17.6%
YouTube Views	N/A	14,739	


CONCLUSION

- CALM achieved visibility in all 3 strategies (mass marketing, digital marketing, and community engagement)
 - Increased website traffic post-launch suggests greater public awareness and engagement.
 - Preliminary analysis indicates a correlation between the CALM campaign and reduced DUP for first-episode psychosis treatment
 - Reduced DUP supply and demand suggest expedited entry into clinics and medication initiation.
 - The study emphasizes the need for a multi-disciplinary approach to psychosis including early detection, timely treatment, and public education.
 - Further research should analyze DUP for patients referred via CALM and survey results from referral sources.
- Current and future directions:
- Developing strategy to track CALM impact longitudinally
 - Analyzing quiz data and implementing 'intervention' to increase those taking the next step to discussing their results
 - How to increase email subscribers
 - Calculating community-level DUP
 - Expanding CALM early detection as we expand EPIC-NOLA clinics across Louisiana.

REFERENCES

Sihari, V. H. et al. (2022). Reducing the Duration of Untreated Psychosis (DUP) in a US Community: A Quasi-Experimental Trial. *Schizophrenia bulletin open*, 3(1), sgab057.





Tulane
University